
- 1) A particle is moving with its position defined by  $s(t) = 2t^3 6t^2 18t + 7$  where t is seconds and s is in feet.
  - a) What are the particle's velocity and acceleration functions?
  - b) What is the total distance traveled by the particle in the first three seconds
  - c) What is the displacement of the particle after the first eight seconds?

$$s(t) = 2t^3 - 6t^3 - 18t + 7$$
 $s(t) = 6t^3 - 12t - 18$ 
 $s(t) = 12t - 12$ 
 $s(0) = 7$ 



- 2) A silver dollar is dropped from a building that is 1,296 feet in height. Time is in second
  - a) What are the silver dollar's height, velocity and acceleration functions?
  - b) When does the silver dollar hit the ground and what is its impact velocity?
  - c) How far does the silver dollar travel between 1 and 2 seconds?

1) 
$$h(t) = -16t^{2} + 0t + 1396$$
 $v(t) = -32t$ 
 $a(t) = -32$ 
 $h(t) = 0$ 
 $v(q) = -32(q)$ 
 $-16t^{2} + 1396 = 0$ 
 $-16t^{2} = -1396$ 
 $t^{2} = 81$ 
 $t^{2} = 81$ 

- 3) The displacement in feet of a body moving along a line at any time t in seconds is given  $s(t) = t^3 7t^2 + 8t + 4$ .
  - a) What are the velocity and acceleration functions?
  - b) Find the total distance traveled in the first 5 seconds. Sketch a motion schemat
  - c) What is the velocity of the body when the position is 8 feet?

$$|S(t) = t^{\frac{3}{2}} 7t^{\frac{3}{2}} + 8t + 4/b) \le 0 = 4$$

$$v(t) = 3t^{\frac{3}{2}} - 19t + 8$$

$$a(t) = 6t - 14$$

$$S(t) = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

$$t^{\frac{3}{2}} - 7t^{\frac{3}{2}} + 8t + 4 = 8$$

- 4) A marble is thrown straight  $\underline{down}$  from the top of a 220-foot building. Its initial veloc was 22 feet per second.
  - a) What are the marble's height, velocity, and acceleration functions?
  - b) When does the marble hit the ground and what is its impact velocity?
  - c) What are the velocity and position at three seconds?
  - d) What is its velocity after falling 108 feet?

i) 
$$h(t) = -16t^2 - 22t + 220$$
  
 $u(t) = -32t - 22$   
b)  $h(t) = 0$   
 $-16t^2 - 22t + 220 = 0$   
 $t = 3.084$  sec  
 $u(3084) = -120.688$  A/Sec

c) 
$$h(3) = 10 + 1$$
  
 $V(3) = -118 + 1/3 = 112$   
 $-16 + 2 - 23 + 1 + 2 = 112$   
 $+ = 2 + 2 = 1$   
 $V(3) = -86 + 1/3 = 112$ 

- 5) An object has its position defined by  $s(t) = t^3 8t^2 + 5t + 2$  in feet. Time is in seconds
  - a) What are the velocity and acceleration functions?
  - b) What is the total distance traveled by the object during the first eight seconds?
  - c) What is the displacement of the object after the first eight seconds?
  - d) What is the position when the velocity is 3.1 feet per second?
  - e) What is the velocity when the acceleration in -2.7 feet per second2?

- 6) A bag of sugar is launched vertically upward from a height of 455 feet with an initial velocity of 102 feet per seconds.
  - a) What are the bag's height, velocity, and acceleration functions?
  - b) What is the position of the bag when the velocity is 6 feet per second?
  - c) When will the bag hit the ground? What is its impact velocity?
  - d) When will the bag reach its maximum height? What is its maximum height?
  - e) What is the velocity of the bag when it is 250 feet above the ground?

$$\begin{array}{lll}
(t) = -16t^{2} + 162t + 455 & c) - 16t^{2} + 162t + 455 = 0 \\
(t) = -32t + 102 & t = 9.45ec. \\
x(t) = -32 & d) - 32t + 102 = 0 & y(9.4) = -198.8ft/sec
\\
y(t) = 6 & t = 3.188 sec & e) - 16t^{2} + 162t + 45 \\
32t + 162 = 6 & 5(3.188) = 6|7.562ft & e) - 16t^{2} + 162t + 45 \\
t = 3sec & y(7.98) = -153
\end{array}$$

$$\begin{array}{ll}
(t) = -16t^{2} + 162t + 45 \\
(t) = -16t^{2} + 162t + 162t + 45 \\
(t) = -16t^{2} + 162t + 45 \\
(t) = -16t^{2} + 162t + 162t + 45 \\
(t) = -16t^{2} + 162t + 45 \\$$