

Use the theoretical data provided in this lab if you were absent when the lab was performed.

Background

In this activity you will calculate the concentration of a solution of hydrochloric acid. This acid is a combination of water as the solvent, and the gaseous compound, HCl, as the solute. The “active ingredient” of an acid solution is the hydronium ion (often referred to as just a “hydrogen ion”). In the presence of water HCl molecules *ionize* to form separate ions in the solution. The hydrogen “ions” actually combine with water molecules to form *hydronium* ions H_3O^+ . The Cl^- ions remain separate as long as water is present. The ions are free to react with other particles they come in contact with.



Many books simply refer to these ions as follows,



If the acid solution has a rich abundance of solute, and relatively few “plain” water molecules, the acid is called concentrated. If there is an abundance of water, and not many of the solute particles, the acid is called dilute

Another way to refer to the “richness” of the acid is by the term *Molarity*, M . The molarity of a solution tells the number of moles of solute *per* liter of total solution.

$$M = \frac{\text{moles of solute}}{\text{liters of solution}}$$

In past labs, we have used two different concentrations of hydrochloric acid. 1M HCl and 6M HCl. 1M HCl means that every 1 liter of the acid solution holds 1 mole of dissolved HCl. 6M HCl means that every 1 liter of solution holds 6 moles of dissolved HCl.

In this Lab...

You will calculate the number of moles of HCl dissolved in a certain volume of acid by reacting the acid with CaCO_3 . This is essentially a double replacement reaction, with a decomposition of one of the product. Stoichiometry skills will be used to process mole information.

Procedures

1. Obtain the mass of a clean, dry 100-250 mL beaker.
2. Place 5-7 CaCO_3 chips in the beaker. Obtain the mass of the beakers and chips.
3. The teacher will tell your group how much acid to use. Measure this amount carefully into a graduated cylinder and add it to the CaCO_3 . Wear your goggles and wipe up spills – the acid may be irritating.
4. Place the beaker above the Bunsen burner – heat the content gently – DO NOT BOIL THE ACID – until all “fizzing has stopped. This is evidence that the HCl has been used up.
5. Rinse the remaining CaCO_3 chips a few times with fresh water, and decant the solution. Do not discard any leftover solid material.
6. Heat the CaCO_3 chips gently until they are dry. Let the beaker cool and mass it.

Data

*The grayed boxes are not to be used.
The white boxes should contain measured and calculated values that pertain to this lab!*

Collected Data		Calculated Data	
Mass of beaker and CaCO ₃ chips	71.11 g		
Mass of beaker (clean, empty, dry)	68.18 g		
		Mass of CaCO ₃ before reaction	g
Mass of beaker and leftover CaCO ₃	69.56 g		
		Mass of CaCO ₃ leftover after the reaction	g
		Mass of CaCO ₃ used up in the reaction	g
Volume of acid used in the reaction	30.0 mL		

Analysis

1. Write the balanced equation.
2. Calculate the number of moles of CaCO₃ **used up** in the reaction.
3. Examine the balanced equation for this reaction and use the coefficients to determine the number of moles of HCl that were used up in your reaction.
Important Note: *You cannot use 1 mol = 22.4 L because that is only used for gases at STP.*
4. Convert the volume of acid used to liters.
5. Calculate the **experimental** molarity of the acid you used.
6. Explain how it is that students who used more (or less) acid than you did will still obtain the same molarity of acid.
7. A measure of the molarity of a solution is more meaningful than saying that a solution is “concentrated” or “dilute.” Explain.
8. Using the actual molarity that was given by the teacher and your experimental molarity, calculate the percent error for this experiment.
9. The molarity (concentration) of your acid is also the concentration of hydrogen ion [H⁺] in solution because hydrochloric acid is a strong acid that dissociated completely. Using this concentration, calculate the pH of the acid solution that was used in this experiment.