

Calculus Stuff To Know Cold

Basic Derivatives

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

$$\frac{d}{dx}(\sin x) = \cos x$$

$$\frac{d}{dx}(\cos x) = -\sin x$$

$$\frac{d}{dx}(\tan x) = \sec^2 x$$

$$\frac{d}{dx}(\sec x) = \sec x \tan x$$

$$\frac{d}{dx}(\cot x) = -\csc^2 x$$

$$\frac{d}{dx}(\csc x) = -\csc x \cot x$$

$$\frac{d}{dx}(\ln x) = \frac{1}{x}$$

$$\frac{d}{dx}(e^x) = e^x$$

The Intermediate Value Theorem

If the function $f(x)$ is continuous on $[a,b]$, then for any number c between $f(a)$ and $f(b)$, there exists a number k in the open interval (a,b) such that $f(k)=c$

The Mean-Value Theorem

If a function $f(x)$ is continuous on $[a,b]$ and the first derivative exists on the interval (a,b) , then there exists a number c such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

Rolle's Theorem

If the function $f(x)$ is continuous on $[a,b]$, the first derivative exists on the interval (a,b) and $f(a)=f(b)$; then there exists a number c on (a,b) such that $f'(c) = 0$

Curve Sketching and Analysis

- Find all x-intercepts and y-intercepts
 - x-intercepts** can be found by letting $y=0$
 - y-intercepts** can be found by letting $x=0$
- Find all asymptotes
 - Vertical Asymptotes** – can be found by letting the denominator of a rational expression equal zero
 - Horizontal Asymptotes**-can be found by finding $\lim_{x \rightarrow \pm\infty} f(x)$
 - Oblique Asymptotes**-can be found using long division
- Find all relative extreme values where $f'(x) = 0$ or is undefined
- Find all points of inflection where $f''(x) = 0$ or is undefined
- Plot all critical points, points of inflection, and intercepts
- Plot any additional points you want

Differentiation Rules

Chain Rule: $\frac{d}{dx}[f(u)] = f'(u) \frac{du}{dx}$

Product Rule: $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$

Quotient Rule: $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{\frac{du}{dx}v - u\frac{dv}{dx}}{v^2}$

Average Value of a Function

If the function $f(x)$ is continuous on $[a,b]$ and the first derivative exists on (a,b) , then there exists a number c in (a,b) such that

$$f(c) = \frac{1}{b-a} \int_a^b f(x) dx$$

This value $f(c)$ is the "average value" of the function on the interval $[a,b]$

The Fundamental Theorem of Calculus

$$\int_a^b f(x) dx = F(b) - F(a)$$

The 2nd Fundamental Theorem of Calculus

$$\frac{d}{dx} \int_{a(x)}^{b(x)} f(t) dt = f[b(x)]b'(x) - f[a(x)]a'(x)$$

More Derivatives

$$\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$$

$$\frac{d}{dx}(\cot^{-1} x) = \frac{-1}{1+x^2}$$

$$\frac{d}{dx}(\sec^{-1} x) = \frac{1}{|x|\sqrt{x^2-1}}$$

$$\frac{d}{dx}(\csc^{-1} x) = \frac{-1}{|x|\sqrt{x^2-1}}$$

Solids of Revolution

Disk Method:

$$V = \pi \int_a^b [R(x)]^2 dx$$

Washer Method:

$$V = \pi \int_a^b [R(x)]^2 - [r(x)]^2 dx$$

Shell Method:

$$V = 2\pi \int_a^b xf(x) dx$$

Cross Sections:

$$V = \int_a^b A(x) dx$$

Distance, Velocity, and Acceleration

Let $s(t)$ be position

Velocity or Rate: $v(t) = s'(t)$

Speed = $|v|$

Acceleration: $a(t) = v'(t) = s''(t)$

Average Velocity over $[a,b] = \frac{f(b) - f(a)}{b - a}$

Total Distance = $\int_a^b |v| dt$

where a is initial time and b is final time.

The Trapezoidal Rule

$$\int_a^b f(x) dx \approx \frac{b-a}{2n} [f(x_0) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(x_n)]$$

Integration by Parts

$$\int u dv = uv - \int v du$$

<p>Simpson's Rule: Must use an even number of subdivisions!</p> $\int_a^b f(x) dx \approx \frac{b-a}{3n} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \dots + 4f(x_{n-1}) + f(x_n)]$		<p>Integral of a Natural Log</p> $\int \ln x dx = x \ln x + C$
<p>Distance, Velocity, Acceleration with Vectors</p> <p>Velocity vector = $\left\langle \frac{dx}{dt}, \frac{dy}{dt} \right\rangle$</p> <p>Speed = $\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$</p> <p>Total Distance = $\int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$</p> <p>where a is initial time and b is final time.</p>	<p>Lengths of Curves</p> <p>Rectangular:</p> $L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$ <p>Parametric:</p> $L = \int_a^b \sqrt{\left(\frac{dy}{dt}\right)^2 + \left(\frac{dx}{dt}\right)^2} dt$ <p>Polar:</p> $L = \int_\alpha^\beta \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$	<p>L'Hopital's Rule</p> <p>If $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{0}{0}$ or $\frac{\infty}{\infty}$, then</p> $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$
<p>Geometric Series:</p> $\sum_{n=1}^{\infty} ar^{n-1} = \sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}$	<p>Average Value:</p> $f_{ave} = \frac{1}{b-a} \int_a^b f(x) dx$	<p>Equation of Tangent Line at x=a</p> $y - f(a) = f'(a)(x - a)$
<p>Taylor Series:</p> <p>If the function f is smooth at x=a, then it can be approximated by the nth degree polynomial</p> $f(x) \approx f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n$		
<p>Maclaurin Series: A Taylor Series about x=0 is called a Maclaurin Series</p> $e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \dots$ $\cos x = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \dots$ $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$ $\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$ $\ln(x+1) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$	<p>The Ratio Test:</p> <p>The series $\sum_{k=0}^{\infty} a_k$ converges if</p> $\lim_{k \rightarrow \infty} \left \frac{a_{k+1}}{a_k} \right < 1$ <p>If limit equals 1, you know nothing.</p>	<p>Polar Curves</p> <p>For a polar curve $r(\theta)$, the area inside a "leaf" is</p> $\int_{\theta_1}^{\theta_2} \frac{1}{2} [r(\theta)]^2 d\theta,$ <p>Where θ_1 and θ_2 are the "first" two times that r=0.</p>
		<p>Convert Polar to Parametric</p> <p>Given $r(\theta)$</p> $x(\theta) = r \cdot \cos \theta$ $y(\theta) = r \cdot \sin \theta$
<p>Lagrange Error</p> <p>If $P_n(x)$ is the nth degree Taylor polynomial of $f(x)$ about c and $f^{(n+1)}(t) \leq M$ for all t between x and c, then</p> $ f(x) - P(x) \leq \frac{M}{(n+1)!} x-c ^{(n+1)}$	<p>Alternating Series Error Bound</p> <p>If $S_N = \sum_{k=1}^N (-1)^k a_k$ is the Nth partial sum of a convergent alternating series, then $S_{\infty} - S_N \leq a_{n+1}$</p>	<p>Euler's Method</p> <p>If given that $\frac{dy}{dx} = f(x, y)$ and that the solution passes through (x_0, y_0) where $y(x_0) = y_0$</p> $y_{new} = y_{old} + dx \cdot \frac{dy}{dx}$